



## Solving Problems with Bounds on Linear Forms in Logarithms

Krish Agrawal\*

Submitted: 16 September 2025 Accepted: 08 October 2025 Publication date: 29 October 2025

DOI: 10.70671/k3gz8m22

**Abstract:** This expository paper explores the theory of linear forms in logarithms and its applications to Diophantine equations. We begin with foundational results on transcendental numbers, including Liouville's theorem and the Gelfond–Schneider theorem, before developing Baker's theory of linear forms in logarithms. The paper concludes with applications to Diophantine equations through the Baker–Davenport method, illustrating these techniques with concrete examples. The purpose of this paper is to provide a step-by-step understanding of this particular area of mathematics, and was written as a part of Euler Circle's Independent Paper and Research Writing Program, in which the author studied the topic independently and wrote this paper over a period of 4 weeks.

**Author keywords:** Linear forms in logarithms; Transcendental number theory; Diophantine equations; Baker's theorem; Number Theory

### Introduction

The study of linear forms in logarithms sits at the intersection of number theory and transcendental number theory. It provides tools to solve equations of the form:

$$\alpha_1^{b_1} \alpha_2^{b_2} \cdots \alpha_n^{b_n} = 1$$

by studying the associated linear form:

$$\Lambda = b_1 \log \alpha_1 + \cdots + b_n \log \alpha_n$$

However, before we explore this form of expression further, we first need to understand the basics of transcendental number theory.

### Algebraic and Transcendental Numbers

#### Basic definitions

**Definition 2.1** (Algebraic Number). *A complex number  $\alpha$  is algebraic if there exists a nonzero polynomial  $P \in \mathbb{Z}[x]$  such that  $P(\alpha) = 0$ . The minimal degree of such a polynomial is called its degree.*

**Example 2.2.** The following are algebraic:

- $\sqrt{2}$  is algebraic of degree 2, since it satisfies  $x^2 - 2 = 0$ .
- The golden ratio  $\phi = \frac{1 + \sqrt{5}}{2}$  is algebraic of degree 2.
- Any rational number  $\frac{p}{q}$  is algebraic of degree 1 via  $qx - p = 0$ .

It is also helpful to note that algebraic numbers are countable, since they arise as roots of polynomials with integer coefficients.

**Definition 2.3 (Transcendental Number).** *A complex number is transcendental if it is not algebraic; that is, it does not satisfy any nonzero polynomial equation with integer coefficients.*

#### Liouville's theorem and constructed transcendents

Joseph Liouville's 1844 theorem was the first major result to establish a criterion for transcendence.

**Theorem 2.4** (Liouville's Approximation Theorem). *For any irrational algebraic number  $\alpha$  of degree  $d \geq 2$ , there exists a constant  $C(\alpha) > 0$  such that for all rational numbers  $\frac{p}{q}$  with  $q > 0$ :*

$$\left| \alpha - \frac{p}{q} \right| > \frac{C(\alpha)}{q^d}$$

*Proof.* Let  $P(x) = a_d x^d + \cdots + a_0 \in \mathbb{Z}[x]$  be the minimal polynomial of  $\alpha$ . By the mean value theorem, for any rational  $\frac{p}{q} \neq \alpha$ :

$$|P(p/q) - P(\alpha)| = |P'(\xi)| \cdot \left| \alpha - \frac{p}{q} \right|$$

for some  $\xi$  between  $\alpha$  and  $p/q$ . Since  $P$  is irreducible,  $P(p/q) \neq 0$  and  $q^d P(p/q) \in \mathbb{Z}$ , so:

$$|P(p/q)| \geq \frac{1}{q^d}$$

\*Corresponding Author: Krish Agrawal. Email: krish.595.2026@doonschool.com  
The Doon School



Siegel's lemma guarantees bounded  $p_{km}$  with:

$$\max |p_{km}| \leq (3N)^{8dN/\log N}$$

Why this bound? The exponential and derivative structure inflates the entries in the system; this bound ensures that a small integer solution still exists.

**Refined Schwarz Lemma:** If  $f$  has  $n$  zeros in  $|z| < R$ , then for  $|z| = r < R$ ,

$$|f(z)| \leq |f(0)| \prod_{k=1}^n \frac{R}{|z_k|} \left(\frac{r}{R}\right)^n \max_{|w|=R} |f(w)|$$

## Extending the zeros

Using the identity  $\alpha^{z+n} = \alpha^z \cdot \alpha^n$ , we obtain:

$$f(\beta + n) = \sum_{k,m} p_{km} \alpha^{k\beta} \alpha^{kn} (\beta + n)^m = \alpha^{n\beta} g(n)$$

where

$$g(z) = \sum_{k,m} p_{km} \alpha^{kz} (z + \beta)^m$$

So  $f(\beta + n) = 0$  for  $n = 0, 1, \dots, N - 1$ .

## Order of vanishing

Let  $s$  be the smallest integer such that  $f^{(s)}(\beta) \neq 0$ . Then,  $s \geq \tau$  and  $f$  have many zeros in a disk around  $\beta$ .

## Maximum modulus estimate

We now use a tool from complex analysis to deduce the upper bounds for the function.

**Maximum Modulus Principle:** If  $f$  is analytic in a domain  $D$  and continuous on  $\bar{D}$ , then  $|f(z)|$  attains its maximum on the boundary  $\partial D$ .

On the circle  $|z| = R = 2N^{1/2}$ :

$$|f(z)| \leq (L+1)^2 \cdot \max |p_{km}| \cdot |\alpha|^{LR} \cdot R^L$$

Substituting:

$$|f(z)| \leq N \cdot (3N)^{8dN/\log N} \cdot |\alpha|^{2N} \cdot (2N^{1/2})^{N^{1/2}}$$

Taking logs:

$$\begin{aligned} \log |f(z)| &\leq \frac{8dN}{\log N} \log(3N) + 2N \log |\alpha| + N^{1/2} \log(2N^{1/2}) \\ &\leq CN \end{aligned}$$

## Schwarz lemma refinement

To complement the global bound obtained via the maximum modulus principle, we invoke a refined Schwarz-type lemma to estimate the derivatives near  $\beta$ . This allows us to bound  $|f^{(s)}(\beta)|$  in terms of the global maximum modulus and the order of vanishing, setting up a contradiction with the algebraic lower bound, which we will see in the subsequent subsection.

We get:

$$\max |f(z)| \leq \left(\frac{R}{N}\right)^{N\cdot\tau} = \left(\frac{2}{N^{1/2}}\right)^{N^2/\log N}$$

So:

$$\log |f(z)| \leq -\frac{N^2}{2} + o(N^2) \Rightarrow |f(z)| \leq e^{-cN^2/\log N}$$

## Lower bound via Diophantine approximation

Since  $\alpha^\beta$  is assumed algebraic and  $f^{(s)}(\beta) \neq 0$ , we apply the following:

**Liouville's Theorem (for algebraic numbers):** Let  $\xi$  be algebraic of degree  $d \geq 2$ . Then, there exists  $C(\xi) > 0$  such that:

$$\left| \xi - \frac{p}{q} \right| > \frac{C(\xi)}{q^d}$$

for all rationals  $p/q$ .

We write  $f^{(s)}(\beta)$  as a polynomial in  $\alpha, \beta$ , and  $\alpha^\beta$ , with integer coefficients bounded using earlier results. Hence,

$$|f^{(s)}(\beta)| \geq (3N)^{-8d^2N}$$

Why this bound? This bound is derived by estimating the minimal polynomial of the algebraic number  $f^{(s)}(\beta)$ . The exponent arises from the number of terms and height estimates in the auxiliary function.

## Cauchy integral bound and contradiction

Using Cauchy's integral formula:

$$f^{(s)}(\beta) = \frac{s!}{2\pi i} \int_{|z|=R} \frac{f(z)}{(z - \beta)^{s+1}} dz$$

Taking absolute values:

$$|f^{(s)}(\beta)| \leq \frac{s! \cdot 2\pi R}{2\pi (R/2)^{s+1}} \cdot \max |f(z)| = \frac{s! \cdot 2^{s+1} \cdot e^{-cN^2/\log N}}{R^s}$$

Now  $s \geq \tau = N/\log N$  implies  $s! \leq e^{DN/\log N}$  and  $R^s \geq (2N^{1/2})^{N/\log N}$ . So we get:

$$|f^{(s)}(\beta)| \leq \frac{e^{DN/\log N} \cdot e^{-cN^2/\log N}}{N^{N/(2\log N)}} = e^{-c'N^2/\log N}$$

But this contradicts the lower bound:

$$(3N)^{-8d^2N} \leq |f^{(s)}(\beta)| \leq e^{-c'N^2/\log N}$$

Taking logs:

$$-8d^2N \log(3N) \leq -\frac{c'N^2}{\log N} \Rightarrow \text{Contradiction for large } N$$

□

## Proof framework

The proof of the Gelfond–Schneider theorem reflects a general template used in transcendence theory:

- Construct an auxiliary function with controlled complexity.
- Impose vanishing conditions at many points.
- Use functional equations to extend the zeros.
- Bound the size of the function from above using complex analysis.
- Establish lower bounds via Diophantine approximation.
- Derive a contradiction between the two bounds.

This methodology has been extended to:

- Values of the exponential function at algebraic points (Hermite–Lindemann).
- Logarithms of algebraic numbers (Gelfond–Schneider).
- Elliptic and abelian functions at algebraic points (Schneider).
- Periods of algebraic varieties (conjecturally).

**Remark 3.3.** *The Gelfond–Schneider theorem resolved part of Hilbert’s 7th problem. It proves that  $2^{\sqrt{2}}$  is transcendental, and that  $e^{\pi i} = -1$  involves transcendentality as well.*

## Baker’s Theory of Linear Forms in Logarithms

<sup>2</sup>

The Gelfond–Schneider theorem represents a major advance in transcendence theory, settling Hilbert’s 7th problem and providing the first general results for the transcendence of values like  $\alpha^\beta$ , where both  $\alpha$  and  $\beta$  are algebraic. However, many problems in number theory involve more complicated expressions—particularly linear combinations of several logarithms of algebraic numbers. A natural question arises: What can be said about expressions such as

$$\Lambda = b_1 \log \alpha_1 + b_2 \log \alpha_2 + \cdots + b_n \log \alpha_n,$$

where  $\alpha_i$  are fixed nonzero algebraic numbers and  $b_i$  are integers? If such a form is nonzero, how small can it be in terms of the sizes of the coefficients and the complexity of the numbers involved?

In the 1960s, Alan Baker developed a powerful generalization of the Gelfond–Schneider theorem, proving that any nontrivial linear form in logarithms of algebraic numbers is not only nonzero but also bounded away from zero by an explicit, effective lower bound. This result, now known as Baker’s theorem, laid the foundation for a large part of effective Diophantine analysis and provided new methods for bounding the size of solutions to exponential equations.

We now state a simplified version of Baker’s theorem, emphasizing its effectiveness and applicability.

**Theorem 4.1** (Baker, 1966). *Let  $\alpha_1, \dots, \alpha_n$  be nonzero algebraic numbers, none equal to 1, and suppose that the logarithms  $\log \alpha_i$  are taken with respect to a fixed branch of the logarithm on  $\mathbb{C} \setminus (-\infty, 0]$ . Let  $b_1, \dots, b_n \in \mathbb{Z}$ , not all zero. Define*

$$\Lambda = b_1 \log \alpha_1 + \cdots + b_n \log \alpha_n.$$

*Then, if  $\Lambda \neq 0$ , we have*

$$|\Lambda| > \exp(-C_1 \cdot \log B \cdot \log A_1 \cdots \log A_n),$$

*where*

- $B = \max\{|b_1|, \dots, |b_n|\}$ ,
- $each A_i \geq \max\{Dh(\alpha_i), |\log \alpha_i|, 0.16\}$ ,
- $h(\alpha_i)$  is the (logarithmic) height of  $\alpha_i$ ,
- $D$  is the degree of the number field  $\mathbb{Q}(\alpha_1, \dots, \alpha_n)$ ,
- and  $C_1$  is an effectively computable constant depending only on  $n$  and  $D$ .

This bound is significant for several reasons. First, it gives a concrete inequality—if one knows the coefficients  $b_i$  and the heights and degrees of the  $\alpha_i$ , then the right-hand side becomes a computable quantity. Second, the bound allows one to conclude that  $\Lambda$  is not too small. In many number-theoretic applications, this means that the expression  $\Lambda$  must actually be bounded away from zero, and therefore an integer linear combination of logarithms cannot vanish.

The logarithmic height  $h(\alpha)$  measures the arithmetic complexity of an algebraic number. For rational numbers  $\alpha = \frac{p}{q}$  in lowest terms, we have  $h(\alpha) = \log \max\{|p|, |q|\}$ . For general algebraic numbers, this extends via consideration of the minimal polynomial and the archimedean absolute values of its conjugates. The appearance of  $h(\alpha_i)$  in the lower bound reflects the intuitive idea that the more complicated the input numbers are, the more flexibility they allow in small linear combinations.

For a full and rigorous proof of Baker’s theorem, the reader is referred to *Transcendental Number Theory* by Alan Baker.<sup>2</sup>

It is also worth emphasizing that Baker’s theorem is effective: the constant  $C_1$  can be computed explicitly given the input data. This is a major departure from earlier transcendence theorems, which often proved existence results without providing any computational information.

The power of this theorem is best appreciated through its applications. For instance, it enables one to solve Diophantine equations involving exponential expressions by first proving that a certain linear form in logarithms is nonzero and then bounding how small it can be. These bounds, though typically extremely small, are nonetheless finite and lead directly to constraints on the size of possible integer solutions.

In the next section, we will see how this theoretical result can be made more practical using a method developed by Baker and Davenport, which allows one to reduce the resulting bounds significantly by incorporating ideas from Diophantine approximation and continued fractions.

## Computing Effective Bounds Using Baker's Theorem

The true power of Baker's theorem lies not just in its qualitative assertion of transcendence or linear independence, but in its capacity to provide explicit lower bounds on expressions involving logarithms of algebraic numbers. This is crucial in many Diophantine contexts, where the goal is to prove that certain equations admit only finitely many solutions, or even to identify those solutions completely.

Suppose we have algebraic numbers  $\alpha_1, \dots, \alpha_n$ , each not equal to 0 or 1, and define the linear form:

$$\Lambda = b_1 \log \alpha_1 + \dots + b_n \log \alpha_n$$

where the coefficients  $b_1, \dots, b_n$  are integers, not all simultaneously zero. We assume  $\Lambda \neq 0$ . The question is: How small can  $|\Lambda|$  be?

Baker's theorem provides a compelling answer. If we let  $B = \max |b_i|$ , let  $h(\alpha_i)$  denote the logarithmic height of  $\alpha_i$ , and let  $D$  be the degree of the number field generated by all  $\alpha_i$ , then there exists a computable constant  $C_1$ , depending only on  $n$  and  $D$ , such that:

$$|\Lambda| > \exp(-C_1 \cdot (\log B)(\log A_1) \cdots (\log A_n))$$

Here, each  $A_i$  is defined as:

$$A_i = \max\{Dh(\alpha_i), |\log \alpha_i|, 0.16\}$$

To see this in action, consider the expression  $\Lambda = x \log 2 - y \log 3$ , which arises in problems like finding solutions to  $2^x = 3^y$ . If  $x, y \leq 1000$ , then  $B = 1000$ , and both 2 and 3 are rational, so  $h(2) = \log 2$ ,  $h(3) = \log 3$ , and  $D = 1$ .

Plugging in, the bound becomes:

$$|\Lambda| > \exp(-C_1 \cdot \log(1000) \cdot \log \log 2 \cdot \log \log 3)$$

With standard estimates for the logarithms and  $C_1$  on the order of  $10^4$ , we find that the lower bound is something like  $10^{-10000}$ . While seemingly negligible, this value is not zero—and that makes all the difference. This bound guarantees a minimum separation between linear combinations of logarithms of algebraic numbers, allowing us to eliminate hypothetical integer solutions with large coordinates.

Although the exponential decay in the bound may feel extreme, especially for large  $B$ , its explicit nature makes it useful in practice. It opens the door to computational refinements such as the Baker–Davenport method, which we now explore.

Despite the strength of Baker's bounds, their numerical size often makes them impractical for direct computation. The Baker–Davenport method addresses this limitation. It combines the theoretical guarantees from Baker's theorem with elementary tools like continued fractions to dramatically reduce the bound and isolate the few possible solutions. This hybrid approach bridges the gap between transcendental number theory and concrete problem-solving.

## Matveev's Refinement of Linear Form Bounds

Baker's theorem provides an effective lower bound for linear forms in logarithms, but the constants involved can be extremely large in practice, limiting the usability of the bounds in computational settings. In 2000, E. M. Matveev introduced a major refinement of these bounds, leading to significantly sharper estimates and better constants for a wide class of problems.<sup>3</sup>

Matveev's theorem applies to linear forms in logarithms of algebraic numbers and provides fully explicit lower bounds that are often several orders of magnitude smaller than those obtained from Baker's original result. These improvements are especially noticeable in applications involving small-degree algebraic numbers or when the logarithmic heights of the numbers are moderate.

In simplified form, Matveev's bound states that for algebraic numbers  $\alpha_1, \dots, \alpha_n \in \overline{\mathbb{Q}}^\times$ , and nonzero integers  $b_1, \dots, b_n$ , if  $\Lambda = b_1 \log \alpha_1 + \dots + b_n \log \alpha_n \neq 0$ , then

$$|\Lambda| > \exp(-C(n) \cdot D^2 \cdot (1 + \log D)(1 + \log B) \cdot (\log A_1) \cdots (\log A_n)),$$

where

- $D$  is the degree of the number field generated by the  $\alpha_i$ ,
- $B = \max\{|b_i|\}$ ,
- $A_i \geq h(\alpha_i)$  are bounds on the logarithmic heights,
- and  $C(n)$  is an explicit constant depending only on  $n$ .

Although this expression resembles Baker's original bound structurally, Matveev's constants are much more favorable. This makes it possible to use the resulting inequalities directly in computations, particularly in bounding solutions to exponential Diophantine equations.

Modern applications often rely on Matveev's version of the bound for practical results. For example, in determining perfect powers in linear recurrence sequences or resolving variants of the Ramanujan–Nagell equation, Matveev's bound can reduce the upper bounds for unknowns from astronomical values to within computational reach.

**Remark 6.1.** For a full proof of Matveev's result and its many variants, the reader is referred to his 2000 paper in the *Izvestiya: Mathematics journal*.<sup>3</sup>

## The Baker–Davenport Reduction Method

While Baker's theorem provides an explicit lower bound on  $|\Lambda|$ , the bound is often so tiny that it alone cannot exclude large integer solutions from consideration. The Baker–Davenport method bridges this gap. It strengthens the application of Baker's bound by combining it with elementary yet powerful tools from Diophantine approximation, particularly continued fractions. This method was introduced in a collaborative paper by Baker and Davenport.<sup>4</sup>

The basic idea is to recast the inequality  $|\Lambda| > \delta$  into a statement about how well a rational number can approximate a certain irrational one. Suppose we have:

$$\Lambda = x \log \alpha - y \log \beta \neq 0$$

We divide both sides by  $x \log \beta$  to obtain:

$$\left| \frac{\log \alpha}{\log \beta} - \frac{y}{x} \right| > \frac{\delta}{|x \log \beta|}$$

This inequality shows that  $\frac{y}{x}$  is not too close to the irrational number  $\log \alpha / \log \beta$ . If we can find rational approximations to  $\log \alpha / \log \beta$  that are closer than the bound allows, we can rule them out as candidates. Conversely, we may use the convergents of the continued fraction expansion of  $\log \alpha / \log \beta$  to approximate this ratio and compare them to known integer solutions.

This technique allows us to zoom in on potential solutions with small values of  $x$  and  $y$ , sharply reducing the upper bounds that arise from Baker's theorem. In this way, what was initially a massive bound, say  $x < 10^{20}$ , can often be brought down to something like  $x < 100$ , making a brute-force search feasible.

Thus, the Baker–Davenport method transforms a theoretical lower bound into a practical algorithm. It is this combination of transcendental number theory and computational approximation that makes the method one of the most effective tools in solving exponential Diophantine equations.

### Worked example: solving $2^x - 3^y = 1$

Consider the equation  $2^x - 3^y = 1$ . This problem asks for pairs of integers  $(x, y)$  such that the difference between a power of two and a power of three equals one. Rearranging, we get:

$$2^x = 3^y + 1$$

Taking logarithms, we find:

$$x \log 2 = \log(3^y + 1)$$

For large  $y$ ,  $3^y + 1$  is very close to  $3^y$ , and so:

$$x \log 2 \approx y \log 3$$

which leads to:

$$\left| \frac{\log 2}{\log 3} - \frac{y}{x} \right| \approx \text{very small}$$

Now, using the continued fraction expansion of  $\log 2 / \log 3 \approx 0.6309$ , we compute convergents such as  $2/3$ ,  $3/5$ ,  $8/13$ , and so on. Each of these gives a candidate rational approximation  $y/x$ . For each such pair  $(x, y)$ , we check whether it satisfies the original equation.

Trying  $x = 1$ , we find:

$$\begin{aligned} 2^1 &= 2, \quad 3^0 = 1, \quad 2 - 1 = 1 \Rightarrow \text{solution found: } (x, y) \\ &= (1, 0) \end{aligned}$$

Trying  $x = 2$ , we find:

$$\begin{aligned} 2^2 &= 4, \quad 3^1 = 3, \quad 4 - 3 = 1 \Rightarrow \text{solution found: } (x, y) \\ &= (2, 1) \end{aligned}$$

Trying  $x = 4$  yields  $2^4 = 16$ , but  $3^2 = 9$  and  $16 - 9 = 7$ , which does not satisfy the equation. As we continue this process, we find that all other convergents violate the bound imposed by Baker's theorem and refined by the continued fraction approximation.

Eventually, this method eliminates all possible solutions except  $(1, 0)$  and  $(2, 1)$ . In this way, the Baker–Davenport method provides a complete and rigorous resolution of the Diophantine equation.

## Applications to Diophantine Equations

### Perfect powers in recurrence sequences

One particularly striking application of linear forms in logarithms arises in the analysis of perfect powers appearing within classical recurrence sequences. A natural example is the Fibonacci sequence  $(F_n)$ , defined by  $F_0 = 0$ ,  $F_1 = 1$ , and the recurrence relation  $F_{n+2} = F_{n+1} + F_n$ .

The question we now pose is whether the Fibonacci sequence contains any perfect squares beyond the trivial examples. Indeed, it is easy to check that  $F_0 = 0$ ,  $F_1 = F_2 = 1$ , and  $F_{12} = 144 = 12^2$ . But are there any others?

To address this, one begins by recalling the closed-form expression for  $F_n$ , known as Binet's formula:

$$F_n = \frac{\phi^n - \psi^n}{\sqrt{5}}$$

where  $\phi = \frac{1 + \sqrt{5}}{2}$  is the golden ratio and  $\psi = \frac{1 - \sqrt{5}}{2}$  is its algebraic conjugate. As  $n$  grows, the term  $\psi^n$  becomes negligible since  $|\psi| < 1$ , so we approximate:

$$F_n \approx \frac{\phi^n}{\sqrt{5}}$$

Now, suppose  $F_n$  is a perfect square, say  $F_n = y^2$  for some integer  $y$ . Then:

$$\frac{\phi^n}{\sqrt{5}} \approx y^2 \quad \Rightarrow \quad \phi^n \approx \sqrt{5}y^2$$

Taking logarithms on both sides, we obtain:

$$n \log \phi \approx \log \sqrt{5} + 2 \log y$$

and hence,

$$|n \log \phi - 2 \log y - \log \sqrt{5}| \approx 0$$

This is now a linear form in logarithms of algebraic numbers:  $n \log \phi - 2 \log y - \log \sqrt{5}$ . Applying Baker's theorem to this expression provides an effective lower bound on its absolute value, which in turn leads to an explicit upper bound on  $n$ . Typically, the bound from Baker's theorem is large, but it can be substantially reduced using the Baker–Davenport method.

Once an upper bound is known, one simply checks all Fibonacci numbers  $F_n$  for  $n$  up to that bound and verifies whether any of them is a perfect square. Such an analysis shows that the only values of  $n$  for which  $F_n$  is a perfect square are  $n = 0, 1, 2, 12$ . No other Fibonacci numbers are perfect squares.

This type of argument, which combines the closed-form representation of a recurrence with bounds on linear forms in logarithms, has proven extremely effective in answering questions about perfect powers in recurrence sequences more generally.

### The Thue equation

Another important application of Baker's theory involves solving Thue equations. These are Diophantine equations of the form:

$$F(x, y) = m$$

where  $F(x, y)$  is an irreducible homogeneous binary form of degree at least three with integer coefficients, and  $m$  is a nonzero integer. The classic result due to Thue tells us that for a fixed  $F$  and  $m$ , this equation has only finitely many integer solutions  $(x, y)$ . However, Thue's original proof is ineffective—it gives no means to compute or bound the solutions.

Baker's theory changes this entirely. It allows us to transform the problem into a system involving linear forms in logarithms, thereby furnishing explicit upper bounds on  $|x|$  and  $|y|$ .

The argument begins by factorizing the binary form  $F(x, y)$  over the algebraic closure of  $\mathbb{Q}$ . Suppose we have:

$$F(x, y) = \prod_{i=1}^d (\alpha_i x - \beta_i y)$$

where the  $\alpha_i, \beta_i$  lie in some finite extension of  $\mathbb{Q}$ , and  $d \geq 3$  is the degree of  $F$ . Now, since  $F(x, y) = m$ , we see that for at least one index  $i$ , the quantity  $|\alpha_i x - \beta_i y|$  must be small—no larger than roughly  $|m|^{1/d}$ .

This gives us an approximation:

$$\left| \alpha_i \frac{x}{y} - \beta_i \right| \ll |y|^{-d}$$

Taking logarithms, we consider expressions of the form:

$$\log \left| \alpha_i \frac{x}{y} - \beta_i \right|$$

These are precisely the kinds of quantities to which Baker's bounds apply. Using the theory of linear forms in logarithms, we can place explicit lower bounds on such logarithmic expressions. Matching this with the upper bounds obtained from the factorization and the size of  $m$ , we deduce explicit upper bounds on  $|x|$  and  $|y|$ .

Once those bounds are known, even if large, they reduce the original infinite problem to a finite one: we simply need to check all integer pairs  $(x, y)$  within the bounded region to find all solutions to the Thue equation.

In practice, further refinements, often using reduction techniques and continued fraction approximations, can decrease the computational load significantly. This makes Baker's theory not only theoretically satisfying but also practically viable for determining the complete set of solutions to many Diophantine equations once thought intractable.

### Conclusion

Linear forms in logarithms form a central part of modern transcendental number theory, providing explicit techniques for dealing with exponential Diophantine equations. What distinguishes this area is not only its ability to prove the transcendence of certain numbers but also its effectiveness in yielding concrete numerical bounds on the size of integer solutions. This paper has shown how classical theorems like those of Liouville and Gelfond–Schneider lay the foundation for Baker's general result. Through Baker's theorem and its computational refinements, especially the Baker–Davenport method, we obtained tools capable of producing meaningful results in practice. The examples examined, from exponential equations such as  $2^x = 3^y + 1$  to the study of perfect powers in recurrence sequences, illustrate how linear forms in logarithms can be used to fully resolve equations that otherwise admit infinitely many possibilities.

Although the bounds provided by Baker's theorem are often too large for direct application, methods like continued fraction approximation allow one to reduce them substantially. This combination of transcendental estimates with classical number-theoretic techniques is what gives the method its strength.

### Acknowledgments

I would like to express my sincerest gratitude to my mentors, Justin Wu and Simon Rubinstein-Salzedo, without whose guidance and support I would not have been able to explore the field of transcendental number theory or pursue this particular topic.

## References

- [1] Gelfond AO. *Transcendental and Algebraic Numbers*. Dover Publications; 1960.
- [2] Baker A. *Transcendental Number Theory*. Cambridge University Press; 1975.
- [3] Matveev EM. An explicit lower bound for a homogeneous rational linear form in logarithms of algebraic numbers. *Izv Math*. 2000;64(6):1217–1269.
- [4] Baker A, Davenport H. The equations  $3x^2 - 2 = y^2$  and  $8x^2 - 7 = z^2$ . *Q J Math*. Oxford Series; 1969;20(2):129–137.
- [5] Lang S. *Introduction to Transcendental Numbers*. Addison-Wesley; 1966.
- [6] Shorey TN, Tijdeman R. Exponential diophantine equations. In: *Cambridge Tracts in Mathematics*. Cambridge University Press; 1986.

## About the Authors



**Krish** is currently a high school senior studying at The Doon School pursuing the IB Diploma, and intends to study Physics in University. His paper is an expository paper exploring Linear Forms in Logarithms, an important result in the field of transcendental number theory.